REPORT DOCUMENTATION PAGE Public reporting burden for this collection of information is estimated to exercise. I hour per reported and information is estimated to exercise.			Form Approved OMB No. 0704-0188	
gathering and maintaining the data needed collection of information, including suggest Davis Highway, Suite 1204, Artington, VA 2 1. AGENCY USE ONLY (Leave L	s, and completing and reviewing the collection for reducing this burden, to Washington 22024302, and to the Office of Management blank) 2. REPORT DATE	ir per response, including the time fi in of information. Send comments in in Headquarters Services, Directorat I and Budget, Paperwork Reduction	or reviewing instructions, searching existing data sources, regarding this burden estimate or any other aspect of this et or information Operations and Reports, 1215 Jefferson Project (0704-0188), Washington, DC 20503.	
	ABINA) 2. REPORT DATE	3. REPORT TYPE	AND DATES COVERED - 9/30/96	1
4. TITLE AND SUBTITLE			5. FUNDING NUMBERS	4
Arsenic in Anaero	d Transformations of obic Sediments	Mercury and	Award No.: N00014-96-1-0036	
6. AUTHOR(S) Francois M. M. Mc	orel			
7. PERFORMING ORGANIZATION	NAME(S) AND ADDRESS(ES)		8. PERFORMING ORGANIZATION	
Department of Geosciences			REPORT NUMBER	
Princeton University Guyot Hall			001-96	ļ
Princeton, New Je	ersey 08544			
9. SPONSORING/MONITORING A	GENCY NAME(S) AND ADDRESS	(ES)	10. SPONSORING / MONITORING	Į
Office of Naval R Ballston Centre T 800 North Quincy Arlington, VA 22	Research, ONR 252: D Cower One Street	iane Gales	AGENCY REPORT NUMBER	
1. SUPPLEMENTARY NOTES			- 10060007	000
	·		19960827	บชเ
				~~
2a. DISTRIBUTION/AVAILABILITY	Y STATEMENT		12b. DISTRIBUTION CODE	Ī
			1	
Approved for publ	ic release distribut	tion unlimited		
Approved for publ 3. ABSTRACT (Maximum 200 wo)		tion unlimited		
The objectives of precipitation and dissolvent consequences of these preamd their accumulation in Based on the preament arsenic and the mercury arsenate to arsenite and the two processes (reduction of arsenic in anaerobic serates of oxidation and repreliminary data, it appears	of this project are to elucidate ution of arsenic and mercu ocesses in determining the polymer are organisms. Eliminary research described aspects of this project. In the precipitation of arsenite as of arsenate and precipitation ediments. The rate of dissolved mercu	e the chemical and biology in anaerobic coasts atential release of these in the proposal, rapid p particular, the condition the arsenic trisulfide so of arsenite) largely detaition of mercuric sulfidery are also being quare proposal, the presence	ogical mechanisms that control the al sediments and to quantify the toxic elements to the water column progress has been made on both the cons for the microbial reduction of colid have been documented. These the ermine the environmental mobility the under various conditions and the intified. In addition, according to the of polysulfides can enhance many	_
The objectives of precipitation and dissolution consequences of these produced and their accumulation in Based on the precipitation and the mercury arsenate to arsenite and the two processes (reduction of arsenic in anaerobic serates of oxidation and repreliminary data, it appear fold the rate of microbial	of this project are to elucidate ution of arsenic and mercu ocesses in determining the post marine organisms. eliminary research described aspects of this project. In the precipitation of arsenite as of arsenate and precipitation ediments. The rate of dissolute duction of dissolved merculars that, as hypothesized in the	e the chemical and biology in anaerobic coasts atential release of these in the proposal, rapid p particular, the condition the arsenic trisulfide so of arsenite) largely detaition of mercuric sulfidery are also being quare proposal, the presence	ogical mechanisms that control the all sediments and to quantify the toxic elements to the water column progress has been made on both the constant for the microbial reduction of colid have been documented. These ermine the environmental mobility is under various conditions and the intified. In addition, according to	-
The objectives of precipitation and dissol- consequences of these pre and their accumulation in Based on the pre arsenic and the mercury arsenate to arsenite and the two processes (reduction of arsenic in anaerobic serates of oxidation and repreliminary data, it appear fold the rate of microbial	of this project are to elucidate ution of arsenic and mercu occesses in determining the polymer organisms. Eliminary research described aspects of this project. In the precipitation of arsenite as of arsenate and precipitation ediments. The rate of dissoluted distribution of dissolved merculars that, as hypothesized in the methylation of mercury in an	e the chemical and biology in anaerobic coasts atential release of these in the proposal, rapid p particular, the condition the arsenic trisulfide so of arsenite) largely detaition of mercuric sulfidery are also being quare proposal, the presence aerobic sediments.	ogical mechanisms that control the al sediments and to quantify the toxic elements to the water column progress has been made on both the cons for the microbial reduction of colid have been documented. These ermine the environmental mobility e under various conditions and the intified. In addition, according to co of polysulfides can enhance many	<u>-</u>
The objectives of precipitation and dissolvent consequences of these preamd their accumulation in Based on the preamsenic and the mercury arsenate to arsenite and the two processes (reduction of arsenic in anaerobic serates of oxidation and repreliminary data, it appears fold the rate of microbial Arsenic, Mercury,	of this project are to elucidate ution of arsenic and mercu ocesses in determining the post marine organisms. eliminary research described aspects of this project. In the precipitation of arsenite as of arsenate and precipitation ediments. The rate of dissolute duction of dissolved merculars that, as hypothesized in the	e the chemical and biology in anaerobic coasts atential release of these in the proposal, rapid p particular, the condition the arsenic trisulfide so of arsenite) largely detaition of mercuric sulfidery are also being quare proposal, the presence aerobic sediments.	ogical mechanisms that control the al sediments and to quantify the toxic elements to the water column progress has been made on both the cons for the microbial reduction of colid have been documented. These ermine the environmental mobility e under various conditions and the intified. In addition, according to co of polysulfides can enhance many	-
The objectives of precipitation and dissol- consequences of these pre and their accumulation in Based on the pre arsenic and the mercury arsenate to arsenite and the two processes (reduction of arsenic in anaerobic serates of oxidation and repreliminary data, it appear fold the rate of microbial	of this project are to elucidate ution of arsenic and mercu occesses in determining the polymer organisms. Eliminary research described aspects of this project. In the precipitation of arsenite as of arsenate and precipitation ediments. The rate of dissoluted distribution of dissolved merculars that, as hypothesized in the methylation of mercury in an	e the chemical and biology in anaerobic coasts atential release of these in the proposal, rapid p particular, the condition the arsenic trisulfide so of arsenite) largely detaition of mercuric sulfidery are also being quare proposal, the presence aerobic sediments.	ogical mechanisms that control the al sediments and to quantify the toxic elements to the water column progress has been made on both the cons for the microbial reduction of colid have been documented. These ermine the environmental mobility is under various conditions and the intified. In addition, according to so of polysulfides can enhance many	-

Standard Form 298 (Rev. 2-89) Prescribed by ANSI Std. 239-18 298-102

GENERAL INSTRUCTIONS FOR COMPLETING SF 298

The Report Documentation Page (RDP) is used in announcing and cataloging reports. It is important that this information be consistent with the rest of the report, particularly the cover and title page. Instructions for filling in each block of the form follow. It is important to stay within the lines to meet optical scanning requirements.

- Block 1. Agency Use Only (Leave blank).
- Block 2. Report Date. Full publication date including day, month, and year, if available (e.g. 1 Jan 88). Must cite at least the year.
- Block 3. Type of Report and Dates Covered. State whether report is interim, final, etc. If applicable, enter inclusive report dates (e.g. 10 Jun 87 30 Jun 88).
- Block 4. <u>Title and Subtitle</u>. A title is taken from the part of the report that provides the most meaningful and complete information. When a report is prepared in more than one volume, repeat the primary title, add volume number, and include subtitle for the specific volume. On classified documents enter the title classification in parentheses.
- Block 5. <u>Funding Numbers</u>. To include contract and grant numbers; may include program element number(s), project number(s), task number(s), and work unit number(s). Use the following labels:

C - Contract G - Grant PR - Project TA - Task

PE - Program Element WU - Work Unit Accession No.

- Block 6. <u>Author(s)</u>. Name(s) of person(s) responsible for writing the report, performing the research, or credited with the content of the report. If editor or compiler, this should follow the name(s).
- Block 7. <u>Performing Organization Name(s) and Address(es)</u>. Self-explanatory.
- Block 8. <u>Performing Organization Report</u>
 <u>Number</u>. Enter the unique alphanumeric report number(s) assigned by the organization performing the report.
- **Block 9.** <u>Sponsoring/Monitoring Agency Name(s)</u> and <u>Address(es)</u>. Self-explanatory.
- Block 10. Sponsoring/Monitoring Agency Report Number. (If known)
- Block 11. Supplementary Notes. Enter information not included elsewhere such as: Prepared in cooperation with...; Trans. of...; To be published in.... When a report is revised, include a statement whether the new report supersedes or supplements the older report.

Block 12a. <u>Distribution/Availability Statement</u>. Denotes public availability or limitations. Cite any availability to the public. Enter additional limitations or special markings in all capitals (e.g. NOFORN, REL, ITAR).

DOD - See DoDD 5230.24, "Distribution Statements on Technical Documents."

DOE - See authorities.

NASA - See Handbook NHB 2200.2.

NTIS - Leave blank.

Block 12b. Distribution Code:

DOD - Leave blank.

DOE - Enter DOE distribution categories from the Standard Distribution for Unclassified Scientific and Technical Reports.

NASA - Leave blank. NTIS - Leave blank.

- Block 13. <u>Abstract</u>. Include a brief (Maximum 200 words) factual summary of the most-significant information contained in the report.
- Block 14. <u>Subject Terms</u>. Keywords or phrases identifying major subjects in the report.
- Block 15. <u>Number of Pages</u>. Enter the total number of pages.
- Block 16. <u>Price Code</u>. Enter appropriate price code (NTIS only).
- Blocks 17. 19. <u>Security Classifications</u>. Self-explanatory. Enter U.S. Security Classification in accordance with U.S. Security Regulations (i.e., UNCLASSIFIED). If form contains classified information, stamp classification on the top and bottom of the page.
- Block 20. <u>Limitation of Abstract</u>. This block must be completed to assign a limitation to the abstract. Enter either UL (unlimited) or SAR (same as report). An entry in this block is necessary if the abstract is to be limited. If blank, the abstract is assumed to be unlimited.

The Chemistry and Transformations of Mercury and Arsenic in

Anaerobic Sediments

Principal Investigator: François M. M. Morel

Department of Geosciences, Princeton University

Princeton, New Jersey 08544

ONR Grant #: N00014-96-1-0036

First Year Report

The objectives of this project are to elucidate the chemical and biological mechanisms that control the precipitation and dissolution of arsenic and mercury in anaerobic coastal sediments and to quantify the consequences of these processes in determining the potential release of these toxic elements to the water column and their accumulation in marine organisms.

Building on the preliminary research described in the proposal to ONR, we have been able to make rapid progress on both the arsenic and the mercury aspects of this project. In particular we have documented the microbial reduction of arsenate to arsenite and studied the conditions for precipitation of arsenite as the arsenic trisulfide solid. These two processes (reduction of arsenate and precipitation of arsenite) largely determine the environmental mobility of arsenic in anaerobic sediments. We have also begun to quantify the rate of dissolution of mercuric sulfide under various conditions and the rates of oxidation and reduction of dissolved mercury. In addition, according to our preliminary data, it appears that, as hypothesized in the proposal, the presence of polysulfides can enhance many fold the rate of microbial methylation of mercury in anaerobic sediments. These achievements are described succinctly below.

Arsenic: Arsenic occurs in the environment principally in two forms, arsenate, As(V), which forms stable precipitates mainly with iron, and arsenite, As(III), which in contrast has been generally considered quite soluble and hence mobile. The chemical and biological transformations between As(V) and As(III) in oxic and anoxic environments are thus of particular interest. Following our previous work with MIT-13 (now Geospirilum arsenophilus), we have

now isolated a second strain, OREX-4 (now *Desulfitobacterium orex*) that grows by dissimilatory reduction of As(V), and characterized both organisms in terms of their metabolic capabilities. One organism can reduce N(V) as well as As(V), while the other can grow on N(VI) as well as As(V). While MIT-13 is gram-negative, OREX-4 is gram-positive and, according to 16S ribosomal RNA sequencing, the two organisms are phylogenetically quite distant. These results, which are being written-up for publication, demonstrate that As(V) reduction is likely widespread among bacterial phyla and common in contaminated anoxic environs.

In addition to reducing As(V), OREX-4 precipitates As(III) as As_2S_3 in S-containing medium. This precipitation occurs, at least in part, intracellularly and under conditions where no chemical precipitation is observed. This finding puts into question the general assumption that reduction of As(V) to As(III) makes As more mobile in the environment. On the contrary, As_2S_3 precipitation may provide a permanent sink for As in anaerobic sediments. A manuscript on this topic is being submitted to Applied and Environmental Microbiology.

Mercury: The major form of mercury in anoxic sediments is mercuric sulfide (HgS), as cinnabar (red) or metacinnabar (black). These minerals are extremely insoluble and a major question relating to the mobility of Hg in anoxic environments is that of the mechanisms and rate of dissolution of HgS. We have begun a systematic study of the kinetics of HgS dissolution examining the effects of such factors as pH, oxygen concentration, light and the presence of organic acids. At present, it is clear that, as we hypothesized, visible light greatly enchances the kinetics of dissolution as does oxygen. The dissolution process is complicated by the possible reduction of Hg(II) (by sulfur or organic species) and rapid re-oxidation of Hg(0) (ultimately by O₂). The rapid oxidation of Hg(0) in the presence of oxygen seems to have been ignored in the environmental literature and, in many aquatic systems, this process should effectively decrease the net rate of Hg(0) volatilization into the atmosphere. We are in the process of quantifying the rates of each of these chemical reactions under conditions typical of natural sediments.

In addition to the difficulties posed by facile redox reactions, the chemistry of mercury in sediments is complicated by the multiple forms of inorganic sulfur. In particular we have

hypothesized that Hg(II) complexes with polysulfides, S_n^{2-} , may facilitate the bacterial uptake and hence the methylation of mercury. Indeed we have found that addition of 0.1mM polysulfide increases the solubility of HgS about 10 fold at pH's 8-10. Further, in preliminary experiments, the same addition of polysulfides to cultures of methylating bacteria grown under fermentating conditions in the presence of HgS increases many fold the rate of Hg methylation. If these results are confirmed, they will constitute a breakthrough in our understanding of the mechanisms of Hg methylation in anaerobic sediments and of the factors that control it.